

Zootaxa 4231 (3): 398 408 http://www.mapress.com/j/zt/

Copyright © 2017 Magnolia Press

Article



https://doi.org/10.11646/zootaxa.4231.3.6

http://zoobank.org/urn:lsid:zoobank.org:pub:30C80B66-2126-4F13-97EA-AD9175DE1B22

A new species of *Manoa* Fittkau (Diptera: Chironomidae), with DNA barcodes from Xianju National Park, Oriental China

XIN QI^{1, 2}, XIN-HUA WANG³, TROND ANDERSEN⁴ & XIAO-LONG LIN^{5, 6}

 ¹College of Life Sciences, Taizhou University, 318000, Taizhou, China. E-mail: qixin0612@tzc.edu.cn
 ²Management Committee of Xianju National Park, Xianju, 317300, Zhejiang, China
 ³College of Life Sciences, Nankai University, 300071, Tianjin, China. E-mail: xhwang@nankai.edu.cn
 ⁴Department of Natural History, University Museum of Bergen, University of Bergen, NO-5020 Bergen, Norway. E-mail: trond.andersen@uib.no
 ⁵Department of Natural History, NTNU University Museum, Norwegian University of Science and Technology, NO-7491, Trondheim, Norway. E-mail: xiaolong.lin@ntnu.no
 ⁶Corresponding author. E-mail: xiaolong.lin@ntnu.no

Abstract

Manoa juensis sp. n. Manoa xian-

Manoa

Key words:

Introduction

The genus Manoa Fittkau, 1963 was erected by Fittkau (1963) based on M. obscura Fittkau, 1963 from Amazonas state, Brazil. Thereafter two additional species were described: *M. tangae* Andersen & Sæther, 1997 from Tanzania and M. pahayokeensis Jacobsen, 2002 from North America and the Dominican Republic (Andersen & Sæther 1997; Jacobsen & Perry 2002; Silva et al. 2015; R. Jacobsen pers. comm.). Manoa belongs to the tribe Pseudochironomini as proposed by Sæther (1977), and has been suspected to be the most plesiomorphic genus within the tribe, dating back to at least mid-Cretaceous (Andersen & Sæther 1997). Originally Pseudochironomini also included the genera Pseudochironomus Malloch, 1915, Riethia Kieffer, 1917, Aedokritus Roback, 1958, Megacentron Freeman, 1961, and Psilochironomus Sublette, 1966, but the latter genus has been treated as a nomen dubium (see, e.g., Spies & Reiss 1996: 90). A recent addition to the tribe is Madachironomus Andersen, 2016 from Madagascar (Andersen 2016). However, the morphology of many members of Pseudochironomini in Sæther's (1977) wide sense is in serious need of revision, the monophyly of this grouping of genera is not firmly established, and is untested due to sampling shortfall in a molecular phylogeny for the Chironomidae (Cranston et al. 2012). For this reason Epler et al. (2013) considered the validity of a tribe Pseudochironomini as so uncertain that they did not separate if from Chironomini in their work. The adults of Manoa are closely related to Riethia, but can be separated from the latter genus by the notched floor of the vagina of female of Manoa (Andersen & Sæther 1977; Jacobsen & Perry 2002; Trivinho-Strixino et al. 2009).

The genus *Pseudochironomus* was incorrectly recorded from Palaearctic China by Wang (2000) based on larvae which later proved to belong to *Rheotanytarsus pentapoda* (Kieffer) (H.Q. Tang pers. comm.). During our recent investigation in the Xianju National Park, China, we collected adults of a *Manoa* species (Figs 1–4) near a small mountain stream in Kuocangshan Nature Reserve. This is thus the first record of the genus *Manoa* and of the tribe Pseudochironomini from the Oriental region. Below we describe and illustrate *M. xianjuensis* **sp. n.** as adult male and female (associated with the male by DNA barcodes). DNA barcoding using a standard fragment of the

mitochondrial cytochrome *c* subunit 1 gene (COI) has proven a successful method for species-level identification in many animal groups including chironomids (Hebert *et al.* 2003, 2004; Ekrem *et al.* 2007; Lin *et al.* 2015; Montagna *et al.* 2016). However, currently there is no DNA barcode of *Manoa* in the reference library. Consequently we generated the standard DNA barcodes of *M. xianjuensis* **sp. n.** and have uploaded them to the Barcode of Life Data Systems (BOLD, http://www.boldsystems.org).

The Xianju National Park is located in the southeastern part of Zhejiang province, China. The park (28°28'14"–28°59'48"N, 120°17'6"–120°55'51"E) is situated at medium to low latitudes of the northern hemisphere, in a region in which a mid-subtropical monsoon climate prevails. The last glacial period had no significant influence on species extinction in the park. Therefore, Xianju National Park provides fully protected habitats and long-term maintenance of the biological diversity. However, the structure and composition of the fauna and flora in the park still remain rather insufficiently known, particularly concerning insects. Previously, only five species of chironomids were recorded from Xianju National Park (Li *et al.* 2013; Lin *et al.* 2012, 2013; Qi *et al.* 2016a, 2016b).

Material and methods

Upon collection the specimens were preserved in 85% ethanol and stored in the dark at 4°C before morphological and molecular analyses. Genomic DNA was extracted from the thorax and head using Oiagen DNA Blood and Tissue Kits at the Department of Natural History, NTNU University Museum, Trondheim, Norway, following the standard protocol. The final elution volume was 100 I. After DNA extraction, the exoskeleton of each specimen was mounted in Euparal on a microscope slide together with the corresponding wings, legs and antennae, following the procedures outlined by Sæther (1969). Morphological terminology follows Sæther (1980), where applicable. Measurements are given as ranges followed by the mean, when four or more specimens were measured, followed by the number of specimens measured (n) in parentheses.

A 658 bp fragment of the COI region was PCR-amplified using the universal primers LCO1490 and HCO2198 (Folmer *et al.* 1994). DNA amplification was carried out in 25 µl reactions using 2.5 µl 10x Takara ExTaq pcr buffer (CL), 2 µl 2.5 mM dNTP mix, 2 µl 25 mM MgCl₂, 0.2 µl Takara Ex Taq HS, 1 µl 10 µM of each primer, 2 µl template DNA and 14.3 µl ddH₂O. Amplification cycles were performed on a Biorad C1000 Thermal Cycler (Bio-Rad, California, USA) and followed a program with an initial denaturation step of 95°C for 5 min, followed by 34 cycles of 94°C for 30 s, 51°C for 30 s, 72°C for 1 min, and 1 final extension at 72°C for 3 min. PCR products were purified using Illustra ExoProStar 1-Step (GE Healthcare Life Sciences, Buckinghamshire, UK) and shipped to MWG Eurofins (Ebersberg, Germany) for bidirectional sequencing using BigDye 3.1 (Applied Biosystems, Foster City, CA, USA) termination. Sequences were assembled and edited using Sequencher 4.8 (Gene Codes Corp., Ann Arbor, Michigan, USA). Sequence information was uploaded to BOLD SYSTEMS along with an image and collateral information for each voucher specimen. Alignment of the sequences was carried out using the Muscle algorithm (Edgar 2004) on amino acids in MEGA 6 (Tamura *et al.* 2013). The pairwise distances using Kimura 2-Parameter (K2P) model were calculated in MEGA 6. The neighbor joining tree was generated using a K2P substitution model with 1000 bootstrap replications and the "pairwise deletion" option for missing data in MEGA 6.

In addition to our own data, we searched for publicly accessible COI barcodes in BOLD belonging to the genera *Pseudochironomus* and *Riethia* that were longer than 300 base pairs and lacked stop codons. The searches were done on January 3, 2017. Hits were combined with our own data and are available through the public dataset "Pseudochironomini (DS-PSEUDOCH)" in BOLD.

The type specimens of *Manoa xianjuensis* **sp. n.** are deposited in the College of Life Sciences, Nankai University, Tianjin, China (BDN) and at the Zoologische Staatssammlung München, Munich, Germany (ZSM).

Manoa xⁱanjuensⁱs Qi & Lin sp. n.

(Figs 1–7)

Type material. Holotype (deposited in BDN): male (BOLD sample ID: XJ17), China, Zhejiang Province, Taizhou,

Xianju, Kuocangshan Nature Reserve, Yukengcun, near stream, 28.63267°N, 120.57606°E, 190 m a.s.l., 24.V.2016, sweep net, X. Qi. Paratypes: 11 males, (BDN no. XJ18–25, XJ33, XJ36, XJ37), 3 females (ZSM no. XJ34, XJ35; BDN no. XJ16), sampling data as for holotype.

Etymology. Named after the type locality. The species epithet is adjectival for the purposes of nomenclature.

Diagnostic characters. The new species conforms to the genus descriptions in Andersen & Sæther (1977) and Jacobsen & Perry (2002). The male of *Manoa xianjuensis* **sp. n.** can be separated from its congeners by the following combination of characters: frontal tubercles absent; AR 1.13–1.33; wing length 1.43–1.50 mm; wing vein R with 6–8 setae, R_1 with 1–3 setae, squama with 1–3 setae; antepronotum with two lateral setae; LR_1 0.96–0.98; pars ventralis and digitus present. The female of *Manoa xianjuensis* **sp. n.** separated from its congeners by the following combination of characters: frontal tubercles absent; wing length 1.58–1.75 mm; R with 10–12 setae, R_1 with 4 setae, R_{4+5} with 11–12 setae; costal extension with 2–3 non-marginal setae; squama with 2 setae; LR_1 0.90–0.93.

Description. Adult male (n = 4, unless otherwise stated). Total length 2.67–3.08, 2.91 mm. Wing length 1.43– 1.50, 1.49 mm. Total length/wing length 1.80–2.00, 1.92. Wing length/length of profemur 1.84–2.08, 1.97. *Coloration* (Figs 1, 2–4). Body pale apricot yellow.

Head (Fig. 5a, b). Antenna with 13 flagellomeres; AR 1.13–1.33, 1.25. Temporal setae 12–16, 14, including 4 inner verticals, 7–12, 9 outer verticals, and 0–2, 1 postorbitals. Clypeus with 5–12, 9 setae. Tentorium 133–138, 135 mm long; 13–20, 17 mm wide. Palpomere lengths (in mm): 28–35, 32; 25–49, 38; 70–78, 75; 83–95, 90; 108–138, 123. Palpomere ratio (5th/3rd) 1.54–1.77, 1.62. Third palpomere with 2–3, 3 lanceolate sensilla clavata.

Wing (Fig. 5c). VR 1.09–1.32, 1.21. Anal lobe reduced. Membrane microtrichia distinctly visible at 100x magnification. Brachiolum with 2–3, 2 setae; R with 6–8, 7 setae; R_1 with 1–3, 2 setae; costal extension 40–60, 50 mm long; with 0–1, 0 non-marginal setae; remaining veins bare. Squama with 1–3, 2 setae.

Thorax (Fig. 6a). Antepronotum with 2 lateral setae. Acrostichals 9–10, 9; dorsocentrals 7–14, 11 in single row; prealars 2–3, 3; supraalar 1. Scutellum with 6 setae.

Legs (Fig. 5d–f). Spur of fore tibia 38–48, 42 mm long; spurs of mid tibia 33–50, 42 and 32–48, 39 mm long; of hind tibia 50–58, 54 and 45–54, 49 mm long. Apical width of fore tibia 38–48, 40 mm; of mid tibia 38–43, 40 mm; of hind tibia 41–50, 46 mm. Lengths (in mm) and proportions of legs as in Table 1.

TABLE 1.

Manoa xianjuensis sp. n.

Hypopygium (Fig. 6b-d). Laterosternite IX with 2–4, 3 setae. Tergite IX with 36–44, 39 setae; anal point absent. Transverse sternapodeme 25–40, 35 mm long, 13–25, 18 mm wide; phallapodeme 85–113, 101 mm long. Gonocoxite 145–163, 152 mm long; with triangular pars ventralis. Superior volsella 63–75, 70 mm long; 7–13, 10 mm wide; dorsal surface with basal microtrichia and 2–3, 2 setae; ventral surface with numerous long microtrichia and 2 apical setae. Digitus 15–30, 19 mm long; with pointed apex. Inferior volsella pediform, 50–58, 53 mm long; with 9–14, 13 long setae. Median volsella vestigial; with 2–3, 2 apical setae. Gonostylus straight; 95–108, 104 mm long; with pointed apex. HR 1.34–1.58, 1.47; HV 2.46–2.86, 2.67.

Adult female (n = 2, unless otherwise stated). Total length 2.92–3.05 mm. Wing length 1.58–1.75 mm. Total



FIGURE 1. Manoa xianjuensis sp. n., u ,



FIGURE 2. Manoa xianjuensis sp. n., u , . . s – .



FIGURE 3. Manoa xianjuensis sp. n., u , . . s 🦟

Head (Fig. 7a, b). Antenna (Fig. 7a) with 6 flagellomeres; AR 0.30–0.32; flagellomere lengths (in mm): 75–88, 65–68, 68–70, 65 (2), 55 (2), 100–110. Temporal setae 15–16, including 3–4 inner verticals, 6–11 outer verticals and 2–5 postorbitals. Clypeus with 10–13 setae. Tentorium 95–135 mm long, 18–19 mm wide. Palpomere lengths (in mm): 33–40, 45–50, 70–78, 88–90, 150 (1). Palpomere ratio (5th/3rd) 1.92 (1). Third palpomere with 2–3 lanceolate sensilla clavata.

Wing. VR 1.20–1.23. Anal lobe reduced. Membrane microtrichia distinctly visible at 100x magnification. Brachiolum with 2 setae; R with 10–12 setae; R_1 with 4 setae; R_{4+5} with 11–12 setae; costal extension 45–65 mm long, with 2–3 non-marginal setae. Squama with 2 setae.

Thorax (Fig. 7c). Antepronotum with 2 lateral setae. Acrostichals 7–9; dorsocentrals 6–11 in single row, prealars 3, supraalar 1. Scutellum with 6 setae.

Legs. Spur of fore tibia 38–40 mm long, spurs of mid tibia 38–43 and 33–40 mm long, of hind tibia 39–50 and 36–45 mm long. Apical width of fore tibia 38–40 mm, of mid tibia 40–43 mm, of hind tibia 38–48 mm. Lengths (in mm) and proportions of legs as in Table 2.

Abdomen. Numbers of setae on tergites I–VIII: 14–20, 18–28, 18–34, 22–38, 28–30, 24–30, 18–38, 6–20. Numbers of setae on sternites I–VIII: 6–8, 4–7, 12–16, 21–30, 22–31, 24–38, 32–40.

Genitalia (Fig. 7d, e). Sternite VIII divided, with 50–55 mm long notched floor (Fig. 7e) under vagina; floor with approximately 32–44 setae. Tergite IX with 22 setae; gonocoxite IX with 1–2 setae. Cercus 78–100 mm long, with 42–48 setae. Seminal capsules 65–68 mm long, including 13–18 mm long, collar-shaped neck, 38–40 mm wide. Spermathecal ducts curved. Notum 75–88 mm long. Dorsomesal lobe 71–73 mm long.



FIGURE 4. Manoa xianjuensis sp. n., u

TABLE 2.		Manoa xianjuensis sp. n.					
		~	1	2	3	4	
1	680 850	690 750	640 690	290 310	220 250	140 170	
2	780 850	710 770	350 400	150 160	110 130	80 90	
3	880 950	780 830	480 520	230 250	200 (2)	100 130	
	5	LR	BV	SV		BR	
1	90 120	0.92 0.93	2.71 2.76	2.14	4 2.32	2.00 3.00	
2	60 80	0.49 0.52	4.28 4.70	4.05	5 4.26	1.77 2.22	
3	80 90	0.62 0.63	3.43 3.51	3.42	2 3.46	1.80 2.00	

ТА	BI	Æ	2.	

Immature stages. Unknown.

DNA barcoding analysis. The neighbor joining tree (Fig. 8) based on available DNA barcodes unequivocally associates the adult males and females of *M. xianjuensis* sp. n., and supports differentiation of the new species from the other barcoded Pseudochironomini. In addition, we found two distinct clades of under the name Riethia stictoptera Kieffer, 1917 with the maximum intraspecific genetic distance 18.1%, which suggests at least one misidentification in the BOLD data.

Biology and ecology. The new species was collected using a sweep-net close to a small mountain stream in Kuocangshan Nature Reserve in subtropical China. We collected M. xianjuensis sp. n. during the rainy season in late May, when the temperature is 15–20°C.





Key to adult males of Manoa Fittkau

1.	Frontal tubercles present; antepronotal and supraalar setae absent. Brazil		
-	Frontal tubercles absent; antepronotal and supraalar setae present		
2.	Pars ventralis and digitus absent; pulvilli and pseudospurs present. North America, Dominican Republic		
-	Pars ventralis and digitus present; pulvilli and pseudospurs absent		
3.	Wing length 1.81–1.91 mm; AR 1.91–2.19; LR ₁ 1.41; wing vein R ₁ bare; transverse sternapodeme with oral projections. Tan-		
	zania		
-	Wing length 1.43–1.50 mm; AR 1.13–1.33; LR ₁ 0.96–0.98; wing vein R ₁ with 1–3 setae; transverse sternapodeme without oral		
	projections. Oriental China		

Key to adult females of Manoa Fittkau

1.	Frontal tubercles present; antepronotal and supraalar setae absent. Brazil
-	Frontal tubercles absent; antepronotal and supraalar setae present
2.	Wing length 1.00–1.16 mm; squama bare. North America
-	Wing length > 1.50 mm; squama setose
3.	Wing length 2.00–2.06 mm; wing veins R with 17–20 setae, R_1 with 7–8 setae, R_{4+5} with 23–29 setae. Tanzania
-	Wing length 1.58–1.75 mm; wing veins R with 4 setae, R_1 with 10–12 setae, R_{4+5} with 11–12 setae. Oriental China



 FIGURE 7. Manoa xianjuensis sp. n.,
 :)
 ;)
 ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;)
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]
 : ;]



Acknowledgements

We are indebted to Martin Spies (Zoologische Staatssammlung München, Munich, Germany) for comments on the identification, and to Chun-Tang Chen (Xianju Forestry Bureau, Zhejiang, China) for taking photos in the field. Thanks to three anonymous reviewers for their input on the manuscript. Financial support from the National Natural Science Foundation of China (31301908, 31272284), the Zhejiang Provincial Natural Science Foundation of China (LY17C040001) and from the Science & Technology Project of Taizhou (1402ky14) are acknowledged with thanks.

References

- A s , T. (2016) Madachironomus, us. Psu. (D-:C, ., C, .,). M s . CHIRONOMUS Journal of Chironomidae Research, 29, 15 26. s/l = 10.5324/ . 0.29.2113
- A s , T. & S , O.A. (1997) F-s . . . Manoa F- u Psu. . S A . . - - - (D- : C - . - : C - . -). Entomologica scandinavica, 28, 311 317. s//. - /10.1163/187631297 00105
- C s., P.S., H , N.B. & M. s, G.E. (2012) A . . u . . C . . . (D.). Systematic Entomology, 37, 172 188.
- E , R.C. (2004) MUSCLE: u s u – u . u u . Nucleic Acids Research,

32, 1792 1797. , T., W- ss, E. & Su, E. (2007) A . s- DNA su - & ss - . - - - DNA Е s. *Molecular Phylogenetics and Evolution*, 43, 530–542. s//... /10.1016/. .2006.11.021 , J.H., E , T. & C s. , P.S. (2013) T . C ... (D- : C ...). H. ... R ... Е K s ... s s Insect Systematics & Evolution, Supplements, 66, 387 556. z. -. C -. . - su - I. Fr u, E. (1963) *Manoa*, r u G u C r. r (Dr) us Archiv für Hydrobiologie, 59, 373 390. F . - s su u - I . - s z. - s. Molecular Marine Biology and Biotechnology, 3, 294 299 Н Proceedings of the Royal Society of London, Biological Sciences, 270, 313 321. , P.D.N., S. , M. ., , T.S. & F & C.M. (2004) I - s . u DNA . s PLoS Н *Biology*, 2 (10), 312. J. s, R.E. & P, S.A. (2002) A s \sim s. Manoa (D \sim : C \sim . \sim). E sN \sim P. Journal of the North American Benthological Society, 21, 314 325. L₇., L₇, L. & W, H. (2013) N s -s . s. Parametriocnemus G. u . C - (D-, C - . .). ZooKeys, 320, 51 62. s// _ /10.3897/z . s.320.4927 L, L, Q, & W, H. (2012) T. s, S. Bryophaenocladius T, , 1934 (D, , C, , ,) . C - . ZooKeys, 208, 51 60. s// . . . /10.3897/z. . s.208.3378 L, .L., Su, E. & E, T. (2015) E, ------s *PLoS ONE*, 10 (9), 0138993. L~, .L., ., . ., L.+, W.B. & W , .H. (2013) A us Compterosmittia S , 1981 (D- : 1. C - . -) . C - . Zootaxa, 3669 (2), 129 138. , М., М т.V., L т. T. & R. ss., B. (2016) I Т. . DNA B . . . А т. М. s M. (D_ : C . . .). *PloS ONE*, 11 (3), 0149673. Q, ., P, J.W., , R.L. & W, , .H. (2016) Dicrotendipes weigiangensis S. . . (D, : C, . . .) . - , C - . Entomotaxonomia, 38 (3), 208 212. Q_{τ} , , R.L., u, L.F. & W , .H. (2016) A s $-s_{\tau}$ us Polypedilum K- (D- : C $-s_{\tau}$) . u N = P, - . Entomotaxonomia, 38 (2), 131 134. , O.A. (1969) S. N -P. . - , D - s - O . - (D - : C - . -). Bulletin of the S Fisheries Research Board of Canada, 107, 1 154. , O.A. (1977) F --C N --S, S. Bulletin of the Fisheries Research Board of Canada, 197, 1 204. S S , O.A. (1980) G. ss (D- : C . . .). Entomologica scandinavica, Su s, 14, 1 51. S-, F.L., W- u, S. & F, B.D. (2015) A -- su . . -- s(D- : C -. -). D. -- R u -. Chironomus Newsletter on Chironomidae Research, 28, 12–19. Su 22, 61 119. T u , K., S , G. P s , D., Frest A. & Ku , S. (2013) MEGA 6: M. u . ur SS __S s. 6.0. Molecular Biology and Evolution, 30, 2725 2729. s// . ~ /10.1093/ . / s197 T --- . - S --- . S., R. u , F.O. & C s. , P.S. (2009) R s --- . Riethia truncatocaudata (E s. 1931) (D --- : C = ..., b = ..., s = ..., u - . s.s. Riethia. Aquatic Insects, 31, 247 259. s//____ /10.1080/01650420902787556

W, .H. (2000) A S S. ~. ~S. C~ (D~). In: H. ~, O. (E.), Late 20th century research on Chironomidae: an anthology from the 13th International Symposium on Chironomidae. S V, A, .629 652.